

DS 4400

Machine Learning and Data Mining I

Alina Oprea
Associate Professor
Khoury College of Computer Science
Northeastern University


September 24 2020

Outline


- Simple linear regression
 - Lab in Python
 - Example of linear regression fit
- Multiple linear regression
 - Vector and matrix gradients
 - Vectorization
 - Derivation of optimal solution
 - Lab in Python

Supervised Learning: Regression


Training



Testing



Interpretation



$$h_{\theta}(x) = \theta_0 + \theta_1 x$$

$$\text{MSE} = \frac{1}{N} \sum_{i=1}^N [h_{\theta}(x_i) - y_i]^2$$

Solution for simple linear regression

- Dataset $x_i \in R, y_i \in R, h_{\theta}(x) = \theta_0 + \theta_1 x$
- $J(\theta) = \frac{1}{N} \sum_{i=1}^N (\theta_0 + \theta_1 x_i - y_i)^2$ **MSE / Loss**

$$\frac{\partial J(\theta)}{\partial \theta_0} = \frac{2}{N} \sum_{i=1}^N (\theta_0 + \theta_1 x_i - y_i) = 0$$

$$\frac{\partial J(\theta)}{\partial \theta_1} = \frac{2}{N} \sum_{i=1}^N x_i (\theta_0 + \theta_1 x_i - y_i) = 0$$

- Solution of min loss

$$\begin{aligned} -\theta_0 &= \bar{y} - \theta_1 \bar{x} \\ -\theta_1 &= \frac{\sum (x_i - \bar{x})(y_i - \bar{y})}{\sum (x_i - \bar{x})^2} \end{aligned}$$

$$\begin{aligned} \bar{x} &= \frac{\sum_{i=1}^N x_i}{N} \\ \bar{y} &= \frac{\sum_{i=1}^N y_i}{N} \end{aligned}$$

Regression vs Correlation

- **Correlation**
 - Find a numerical value expressing the relationship between variables
- **Regression**
 - Estimate values of response variable on the basis of the values of predictor variable
- The slope of linear regression is related to correlation coefficient
- Regression scales to more than 2 variables, but correlation does not

Lab Simple Linear Regression

```
#!/usr/bin/env python

import numpy as np
import matplotlib.pyplot as plt

import pandas as pd
import seaborn as sns

from sklearn.model_selection import train_test_split
from sklearn.linear_model import LinearRegression
from sklearn.metrics import mean_squared_error

from sklearn.linear_model import Ridge, RidgeCV, Lasso, LassoCV

from sklearn.datasets import load_boston
boston_dataset = load_boston()
```

Boston house prediction dataset

Lab

```
boston = pd.DataFrame(boston_dataset.data, columns=boston_dataset.feature_names)
boston['MEDV'] = boston_dataset.target
boston.head(5)
```

	CRIM	ZN	INDUS	CHAS	NOX	RM	AGE	DIS	RAD	TAX	PTRATIO	B	LSTAT	MEDV
0	0.00632	18.0	2.31	0.0	0.538	6.575	65.2	4.0900	1.0	296.0	15.3	396.90	4.98	24.0
1	0.02731	0.0	7.07	0.0	0.469	6.421	78.9	4.9671	2.0	242.0	17.8	396.90	9.14	21.6
2	0.02729	0.0	7.07	0.0	0.469	7.185	61.1	4.9671	2.0	242.0	17.8	392.83	4.03	34.7
3	0.03237	0.0	2.18	0.0	0.458	6.998	45.8	6.0622	3.0	222.0	18.7	394.63	2.94	33.4
4	0.06905	0.0	2.18	0.0	0.458	7.147	54.2	6.0622	3.0	222.0	18.7	396.90	5.33	36.2

```
print(len(boston))
boston.shape
```

506

```
(506, 14)
```

```

: correlation_matrix = boston.corr().round(2)
  # annot = True to print the values inside the square
  sns.heatmap(data=correlation_matrix, annot=True)

```

: <AxesSubplot:>



Lab

```
# Simple LR
X = pd.DataFrame(np.c_[boston['RM']], columns = ['RM'])

Y = boston['MEDV']

X_train, X_test, Y_train, Y_test = train_test_split(X, Y, test_size = 0.2, random_state=5)
print(X_train.shape)
print(X_test.shape)
print(Y_train.shape)
print(Y_test.shape)
```

```
(404, 1)
(102, 1)
(404,)
(102,)
```

```
slr = LinearRegression()
slr.fit(X_train, Y_train)
```

Lab

```
print(slr.intercept_)  
print(slr.coef_)
```

```
-32.839129906011266  
[8.82345634]
```


```
Y_train_predict = slr.predict(X_train)  
mse = mean_squared_error(Y_train, Y_train_predict)  
  
print("The model performance for training set")  
print('MSE is {}'.format(mse))
```

```
The model performance for training set  
MSE is 48.612648648611334
```

```
df = pd.DataFrame({'Actual': Y_train, 'Predicted': Y_train_predict})  
df.head()
```

	Actual	Predicted
33	13.1	17.463395
283	50.0	37.069115
418	8.8	19.722200
502	20.6	21.160423
402	12.1	23.666285


```
plt.scatter(X_train, Y_train, color='blue')  
plt.plot(X_train, Y_train_predict, color='red', linewidth=2)  
plt.xlabel('RM')  
plt.ylabel('MEDV')  
plt.show()
```



Example Linear Regression Fit

- Consider the following values of predictor and response
 - $X = (-1, 1, 3)$
 - $Y = (-1, 1, 6)$
- Compute $\text{Cov}(X, Y)$ and $\text{Var}[X]$
- Compute $\theta_1 = \frac{\sum (x_i - \bar{x})(y_i - \bar{y})}{\sum (x_i - \bar{x})^2} = \frac{\text{Cov}(X, Y)}{\text{Var}[X]}$
- Compute $\theta_0 = \bar{y} - \theta_1 \bar{x}$
- Find the Least Square Linear Regression model that fits this data optimally

Multiple Linear Regression



- Linear Regression with 2 predictors
- Dataset: $x_i \in R^d, y_i \in R$

Vector Norms

Vector norms: A norm of a vector $\|x\|$ is informally a measure of the “length” of the vector.

$$\|x\|_p = \left(\sum_{i=1}^n |x_i|^p \right)^{1/p}$$

– Common norms: L_1 , L_2 (Euclidean)

$$\|x\|_1 = \sum_{i=1}^n |x_i| \quad \|x\|_2 = \sqrt{\sum_{i=1}^n x_i^2}$$

– L_∞

$$\|x\|_\infty = \max_i |x_i|$$

Vector products

We will use lower case letters for vectors

The elements are referred by x_i .

- Vector dot (inner) product:

$$x^T y \in \mathbb{R} = \begin{bmatrix} x_1 & x_2 & \cdots & x_n \end{bmatrix} \begin{bmatrix} y_1 \\ x_2 \\ \vdots \\ y_n \end{bmatrix} = \sum_{i=1}^n x_i y_i.$$

- Vector outer product:

$$xy^T \in \mathbb{R}^{m \times n} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_m \end{bmatrix} \begin{bmatrix} y_1 & y_2 & \cdots & y_n \end{bmatrix} = \begin{bmatrix} x_1 y_1 & x_1 y_2 & \cdots & x_1 y_n \\ x_2 y_1 & x_2 y_2 & \cdots & x_2 y_n \\ \vdots & \vdots & \ddots & \vdots \\ x_m y_1 & x_m y_2 & \cdots & x_m y_n \end{bmatrix}$$

Hypothesis Multiple LR

- Linear Model

- Consider our model:

$$h(\mathbf{x}) = \sum_{j=0}^d \theta_j x_j$$

- Let

$$\boldsymbol{\theta} = \begin{bmatrix} \theta_0 \\ \theta_1 \\ \vdots \\ \theta_d \end{bmatrix} \quad \mathbf{x}^\top = \begin{bmatrix} 1 & x_1 & \dots & x_d \end{bmatrix}$$

- Can write the model in vectorized form as $h(\mathbf{x}) = \boldsymbol{\theta}^\top \mathbf{x}$

Vector inner product

Training data

	Feature 1		Feature d		
$X =$	1	x_{11}	$\dots$	x_{1d}	Training example i
	$\vdots$	$\vdots$	$\ddots$	$\vdots$	
	1	x_{i1}	$\dots$	x_{id}	
	$\vdots$	$\vdots$	$\ddots$	$\vdots$	
	1	x_{N1}	$\dots$	x_{Nd}	

- Total number of training example: N
- Dimension of training data point (number of features): d
- Dimension of matrix: $N \times (d+1)$

Use Vectorization

- Consider our model for n instances:

$$h(x_i) = \sum_{j=0}^d \theta_j x_{ij}$$

- Let

$$\theta = \begin{bmatrix} \theta_0 \\ \theta_1 \\ \vdots \\ \theta_d \end{bmatrix} \quad \mathbf{X} = \begin{bmatrix} 1 & x_{11} & \dots & x_{1d} \\ \vdots & \vdots & \ddots & \vdots \\ 1 & x_{i1} & \dots & x_{id} \\ \vdots & \vdots & \ddots & \vdots \\ 1 & x_{N1} & \dots & x_{Nd} \end{bmatrix}$$

$\mathbb{R}^{(d+1) \times 1}$
 $\mathbb{R}^{n \times (d+1)}$

Model
parameter

Training
data

- Can write the model in vectorized form as $h_{\theta}(x) = \mathbf{X}\theta$

Model prediction vector $\hat{y}$

Loss function MSE

- For the linear regression cost function:

$$J(\theta) = \frac{1}{N} \sum_{i=1}^N [h_{\theta}(x_i) - y_i]^2$$

$$= \frac{1}{N} \sum_{i=1}^N [\hat{y}_i - y_i]^2$$

$$y = \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ \vdots \\ \vdots \\ y_N \end{bmatrix}$$

$$= \frac{1}{N} ||\hat{y} - y||^2$$
$$= \frac{1}{N} ||X\theta - y||^2$$

$$\hat{y} = \begin{bmatrix} \hat{y}_1 \\ \hat{y}_2 \\ \vdots \\ \vdots \\ \vdots \\ \hat{y}_N \end{bmatrix}$$

Matrix and vector gradients

If $y = f(x)$, $y \in R$ scalar, $x \in R^n$ vector

$$\frac{\partial y}{\partial x} = \left[\frac{\partial y}{\partial x_1} \quad \frac{\partial y}{\partial x_2} \quad \cdots \quad \frac{\partial y}{\partial x_n} \right]$$

Vector gradient
(row vector)

If $y = f(x)$, $y \in R^m$, $x \in R^n$

$$\frac{\partial \mathbf{y}}{\partial \mathbf{x}} = \begin{bmatrix} \frac{\partial y_1}{\partial x_1} & \frac{\partial y_1}{\partial x_2} & \cdots & \frac{\partial y_1}{\partial x_n} \\ \frac{\partial y_2}{\partial x_1} & \frac{\partial y_2}{\partial x_2} & \cdots & \frac{\partial y_2}{\partial x_n} \\ \vdots & \vdots & & \vdots \\ \frac{\partial y_m}{\partial x_1} & \frac{\partial y_m}{\partial x_2} & \cdots & \frac{\partial y_m}{\partial x_n} \end{bmatrix}$$

Jacobian
matrix
(Matrix
gradient)

Properties

- If w, x are $(d \times 1)$ vectors, $\frac{\partial w^T x}{\partial x} = w^T$
- If $A: (n \times d) \ x: (d \times 1)$, $\frac{\partial Ax}{\partial x} = A$
- If $A: (d \times d) \ x: (d \times 1)$, $\frac{\partial x^T Ax}{\partial x} = (A + A^T)x$
- If A symmetric: $\frac{\partial x^T Ax}{\partial x} = 2Ax$
- If $x: (d \times 1)$, $\frac{\partial ||x||^2}{\partial x} = 2x^T$

Min loss function

- Notice that the solution is when $\frac{\partial}{\partial \theta} J(\theta) = 0$

$$J(\theta) = \frac{1}{N} ||X\theta - y||^2$$

Using chain rule

$$f(\theta) = h(g(\theta)), \frac{\partial f(\theta)}{\partial \theta} = \frac{\partial h(g(\theta))}{\partial \theta} \frac{\partial g(\theta)}{\partial \theta}$$

$$h(x) = ||x||^2, g(\theta) = X\theta - y$$

$$\frac{\partial J(\theta)}{\partial \theta} = \frac{2}{N} [(X\theta - y)^T X] = 0 \Rightarrow X^T (X\theta - y) = 0$$

$$(X^T X)\theta = X^T y$$

Closed Form Solution:

$$\theta = (X^T X)^{-1} X^T y$$

Multiple Linear Regression

- Dataset: $x_i \in R^d, y_i \in R$
- Hypothesis $h_\theta(x) = \theta^T x$
- $MSE = \frac{1}{N} \sum (\theta^T x_i - y_i)^2$ Loss / cost

$$\theta = (X^T X)^{-1} X^T y$$

