

Homework 03

Assigned: Tue 04 Oct 2005

Due: Fri 14 Oct 2005

Instructions:

- Feel free to work with others on this assignment. However, you must acknowledge with whom you worked, and you must write up your own solutions.

Problem 1 [15 pts]: Cover and Thomas, Chapter 2, Problems 2, 5, and 6.

Clarification: For each part of Problem 2, state that $H(Y) \sim H(X)$ where \sim is replaced with one of $\{<, \leq, =, \geq, >\}$ and justify your answer.

Problem 2 [30 pts]: Jensen-Shannon Divergence

Let p_1 and p_2 be probability distributions over a discrete space \mathcal{X} and define the *average* of these distributions, \bar{p}_{12} , as follows:

$$\bar{p}_{12}(x) = \frac{p_1(x) + p_2(x)}{2} \quad \forall x \in \mathcal{X}.$$

When the underlying distributions are clear, we shall drop the subscripts; hence, $\bar{p} = \bar{p}_{12}$.

The Jensen-Shannon divergence between two distributions p_1 and p_2 is defined as follows:

$$\text{JS}(p_1, p_2) = \frac{D(p_1 \parallel \bar{p}) + D(p_2 \parallel \bar{p})}{2}.$$

In other words, the Jensen-Shannon divergence is the *average* of the KL-distances to the *average* distribution.

i. Show that

$$\text{JS}(p_1, p_2) = H(\bar{p}) - \frac{H(p_1) + H(p_2)}{2}.$$

In other words, the Jensen-Shannon divergence is the entropy of the *average* minus the *average* of the entropies.

ii. Show that

- $\text{JS}(p_1, p_2) \geq 0$,
- $\text{JS}(p_1, p_2) = \text{JS}(p_2, p_1)$, and
- $\text{JS}(p_1, p_2) = 0$ if and only if $p_1 = p_2$.

These are three of the four properties necessary for a metric, the fourth property being triangle inequality. Additionally, show that

- $\text{JS}(p_1, p_2) \leq 1$.

Hint: To prove this last property, argue that $D(p_i \parallel \bar{p}) \leq 1$ for both $i = 1$ and 2 , then appeal to the definition of Jensen-Shannon.

iii. Let p_1 , p_2 , and p_3 be distributions over a discrete space \mathcal{X} . For Jensen-Shannon to be a metric, it must satisfy the triangle inequality property

$$\text{JS}(p_1, p_2) + \text{JS}(p_2, p_3) \geq \text{JS}(p_1, p_3)$$

in addition to the first three properties described in part (ii) above.

- Show that Jensen-Shannon satisfies the triangle inequality if and only if

$$H(\bar{p}_{12}) + H(\bar{p}_{23}) \geq H(\bar{p}_{13}) + H(p_2).$$

Hint: Use the result from part (i) above.

- Use the above result to show that Jensen-Shannon is *not* a metric by constructing three simple distributions p_1 , p_2 , and p_3 for which the above inequality does not hold. (Distributions over a discrete space of size two suffice.)

Aside: While the Jensen-Shannon divergence is not a metric, it can be shown that the square root of the Jensen-Shannon divergence is a metric. The Jensen-Shannon divergence can also be generalized to allow for weighted averages among distributions and to generate a divergence for an arbitrary number of distributions. The most general form of Jensen-Shannon is as follows: Let p_1, p_2, \dots, p_n be n distributions over a discrete space \mathcal{X} , and let $\vec{\lambda} = \{\lambda_1, \lambda_2, \dots, \lambda_n\}$ be a distribution over $\{1, 2, \dots, n\}$. Define the weighted average distribution $\bar{p}(x) = \sum_i \lambda_i p_i(x) \forall x \in \mathcal{X}$. Then the generalized Jensen-Shannon divergence among p_1, p_2, \dots, p_n with respect to $\vec{\lambda}$ is

$$\begin{aligned} \text{JS}(p_1, p_2, \dots, p_n) &= \sum_i \lambda_i D(p_i \| \bar{p}) \\ &= H(\bar{p}) - \sum_i \lambda_i H(p_i). \end{aligned}$$

Problem 3 [25 pts]: Cover and Thomas, Chapter 2, Problem 15.

Hint: Construct the Venn diagram for intuition.

Problem 4 [30 pts]: Cover and Thomas, Chapter 2, Problem 30.

Hint: This is a constrained optimization problem that you should solve using Lagrange multipliers. You will have two constraints that need to be simultaneously satisfied: one which ensures that the probabilities form a distribution and a second which ensures that the distribution has a specific mean.

Hint: You should obtain $H(X) = (1 + A) \lg(1 + A) - A \lg A$.